



STUDY OF FRACTIONAL INTEGRATION OF THE GENERALIZED MATHIEU SERIES

VARSHA PARIHAR and CHANDRAPAL SINGH CHOUHAN

Research Scholar
Department of Mathematics
and Statistics Bhupal Nobles' University
Udaipur (Raj.)

Supervisor
Department of Mathematics and Statistics
Bhupal Nobles' University, Udaipur (Raj.)

Abstract

In this paper we present some formulas for the Saigo hypergeometric fractional integral involving the generalized Mathieu series $S_{\mu}(r)$, which are expressed in terms of the Hadamard product of the generalized Mathieu series $S_{\mu}(r)$ and the Fox-Wright function $\Psi_q^p(z)$. Corresponding assertions for the classical Riemann-Liouville and Erdélyi-Kober fractional integral deduced. Further, it is emphasized that the results presented here, which are for a seemingly complicated series, can reveal their involved properties via the series of the two known functions.

1. Introduction and Preliminaries

Fractional calculus, which has a long history, is an important branch of mathematical analysis (calculus) where differentiations and integrations can be of arbitrary non-integer order. During the past four decades or so, fractional calculus has been widely and extensively investigated and has gained importance and popularity due mainly to its demonstrated applications in numerous and diverse fields of science and engineering such as turbulence and fluid dynamics, stochastic dynamical system, plasma physics and controlled thermonuclear fusion, nonlinear control theory, image

2020 Mathematics Subject Classification: 18E35.

Keywords: Mathieu series; generalized Mathieu series; fractional calculus operators.

Received March 4, 2022; Accepted September 7, 2022

processing, nonlinear biological systems, and astrophysics. For further details we refer to [1-5].

Now, we recall Saigo fractional integral involving Gauss's Hypergeometric function F_{12} as a kernel. Let $\alpha, \beta, \eta \in C$, $\operatorname{Re}(\alpha) > 0$ and $x > 0$, then Saigo's fractional integral $(I_{0+}^{\alpha, \beta, \eta} f)(x)$, $(I_{0-}^{\alpha, \beta, \eta} f)$ and defined as,

$$(I_{0+}^{\alpha, \beta, \eta} f)(x) = \frac{x^{-\alpha-\beta}}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} F_1^2(\alpha + \beta, -\eta; \alpha; 1 - \frac{t}{x}) f(t) dt \quad (1.1)$$

$$(I_{0-}^{\alpha, \beta, \eta} f)(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} t^{-\alpha-\beta} F_1^2(\alpha + \beta, -\eta; \alpha; 1 - \frac{t}{x}) f(t) dt \quad (1.2)$$

Here and in what follows, $[x]$ denotes the greatest integer less than or equal to the real number x . When $\beta = -\alpha$ the operators in (1.1), (1.2) coincide with the classical Riemann-Liouville fractional integrals and derivatives of order $\alpha \in C$ with $\operatorname{Re}(\alpha) > 0$ and $x > 0$ (see, e.g., [1,4]):

$$(I_{0+}^{\alpha, -\alpha, \eta} f)(x) = (I_{0+}^{\alpha} f)(x) \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} f(t) dt \quad (1.3)$$

$$(I_{0-}^{\alpha, -\alpha, \eta} f)(x) = (I_{0-}^{\alpha} f)(x) = \frac{1}{\Gamma(\alpha)} \int_0^{\infty} (x-t)^{\alpha-1} f(t) dt \quad (1.4)$$

Here and in the following, let C , R^+ , and N be the sets of complex numbers, positive real numbers, and positive integers, respectively, and let $N_0 := N \cup \{0\}$. If $\beta = 0$ in (1.1),(1.2) yields the so-called Erdélyi-Kober fractional integrals of order $\alpha \in C$ with $\operatorname{Re}(\alpha) > 0$ and $x > 0$,

$$(I_{0+}^{\alpha, 0, \eta} f)(x) = (I_{\eta, \alpha}^{\alpha} f)(x) = \frac{x^{-\alpha-\eta}}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} t^{\eta} f(t) dt \quad (1.5)$$

$$(I_{0-}^{\alpha, 0, \eta} f)(x) = (K_{\eta, \alpha}^{-} f)(x) = \frac{x^{\eta}}{\Gamma(\alpha)} \int_0^{\infty} (x-t)^{\alpha-1} t^{-\alpha-\eta} f(t) dt \quad (1.6)$$

A detailed account of such operators along with their properties and applications has been considered by several authors. The following familiar

infinite series of the form

$$S(r) = \sum_{n \geq 1} \frac{2n}{(n^2 + r^2)^2} \quad (1.7)$$

Where $r > 0$, is known in literature as the Mathieu series. Émile Leonard Mathieu was the first to investigate such a series in 1890 in his book elasticity of solid bodies [7]. An alternative version of (7)

$$S^{\sim}(r) = \sum_{n \geq 1} (-1)^{n-1} \frac{2n}{(n^2 + r^2)^2} \quad (1.8)$$

Where $r > 0$, was introduced by Pogány et al. [8]. Closed form integral representations for $S(r)$ and $S^{\sim}(r)$ are given by (see e.g., [8, 9])

$$S(r) = \frac{1}{r} \int_0^{\infty} \frac{x \sin(rx)}{e^x - 1} dx \quad (1.9)$$

and

$$S^{\sim}(r) = \frac{1}{r} \int_0^{\infty} \frac{x \sin(rx)}{e^x - 1} dx \quad (1.10)$$

respectively. Several interesting problems and solutions deal with integral representations and bounds for the following mild generalization of the Mathieu series and its alternative version with a fractional power defined by ([10], p. 2, Equation (16)) (see also, [11], p. 181).

$$S_{\mu}(r) = \sum_{n \geq 1} \frac{2n}{(n^2 + r^2)^{\mu+1}} \quad (1.11)$$

And

$$S_{\mu}^{\sim}(r) = \sum_{n \geq 1} (-1)^{n-1} \frac{2n}{(n^2 + r^2)^{\mu+1}} \quad (1.12)$$

Where $r > 0$, $\mu > 0$, respectively. Such a series has been widely considered in mathematical literature (see, e.g., papers of Cerone and Lenard [10], Diananda [12] and Pogány et al. [8]). Various applications of the familiar Mathieu series and its generalizations in probability theory with other

variants such as trigonometric Mathieu series, harmonic Mathieu series, Fourier-Mathieu series and some other particular forms of the Mathieu series can be found in a recent paper [13]. Recently, Tomovski and Pogány [14] studied the several integral representations of the generalized fractional order Mathieu-type power series

$$S_{\mu}(r; z) = \sum_{n \geq 1} \frac{2nz^n}{(n^2 + r^2)^{\mu+1}} \quad (1.13)$$

Where $r \in R$, $\mu > 0$, $|z| < 1$. Obviously, we have $S_{\mu}(r; 1) = S_{\mu}(r)$ and $S_{\mu}(r; -1) = S_{\mu}^{\sim}(r)$. Various other investigations and generalizations of the Mathieu series with its alternative variants can also be found in [11, 14-24], and the references cited therein. The concept of the Hadamard product (or the convolution) of two analytic functions is useful in our present investigation. It can help us to decompose a newly emerged function into two known functions. If, in particular, one of the power series defines an entire function, then the Hadamard product series defines an entire function, too. Let $f(z) := \sum_{n=0}^{\infty} a_n z^n$, ($|z| < R_f$) and $g(z) := \sum_{n=0}^{\infty} b_n z^n$, ($|z| < R_g$) be two power series whose radii of convergence are denoted by R_f and R_g , respectively. Then, their Hadamard product is the power series defined by

$$(f * g)(z) := \sum_{n=0}^{\infty} a_n b_n z^n = (g * f)(z) \quad (1.14)$$

$|z| < R_g$, where

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n b_n}{a_{n+1} b_{n+1}} \right| = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| \cdot \lim_{n \rightarrow \infty} \left| \frac{b_n}{b_{n+1}} \right| = R_f \cdot R_g.$$

Therefore, in general, we have $R > R_f \cdot R_g$ [25, 26]. For various other investigations involving the Hadamard product (or the convolution), the interested reader may be referred to several recent paper son the subject (see, for example, [27, 28] and the references cited in each of these papers). In this paper, our aim is to study the compositions of the generalized fractional (1)-(2) with the generalized Mathieu series (1.13) in terms of the Hadamard

product (1.14) of the generalized Mathieu series and the Fox–Wright function. Further, corresponding assertions for the classical Riemann–Liouville and Erdélyi–Kober fractional integral are deduced. The results presented in Theorems together with Corollaries are sure to be new and potentially useful, mainly because they are expressed in terms of the Hadamard product with two known functions. At least, a seemingly complicated resulting series expressed in terms of two known functions means that certain properties involved in the complicated resulting series can be revealed via the series of the known functions.

2. Fractional Integration of the Mathieu Series We first recall the Fox–Wright function $\Psi_q^p(z)$ ($p, q \in N_0$) with p numerator and q denominator parameters defined for $\alpha_1, \dots, \alpha_p \in C$ and $b_1, \dots, b_p \in C/Z_0^-$ by (see, for details, [1,3]; see also [4,29]):

$$\begin{aligned} & \Psi_q^p \left[\begin{matrix} (\alpha_1, A_1), (\alpha_2, A_2) \dots (\alpha_p, A_p) \\ (\beta_1, B_1), (\beta_2, B_2) \dots (\beta_q, B_q) \end{matrix} ; z \right] \\ &= \sum_{n=0}^{\infty} \frac{\Gamma(\alpha_1 + A_1 n) \Gamma(\alpha_2 + A_2 n) \dots \Gamma(\alpha_p + A_p n) z^n}{\Gamma(\beta_1 + B_1 n) \Gamma(\beta_2 + B_2 n) \dots \Gamma(\beta_q + B_q n) n!} \end{aligned} \quad (2.1)$$

Where $(A_j \in R^+ (j = 1, 2, 3, \dots, p); B_j \in R^+ (j = 1, 2, 3, \dots, q); 1 + \sum_{j=1}^q B_j - \sum_{j=1}^p A_j \geq 0)$. In particular, when $A_j = B_k = 1 (j = 1, \dots, p; k = 1, \dots, q)$, (23) reduces immediately to the generalized hypergeometric function

$F_q^p(p, q \in N_0)$ (see, e.g., [29]):

$$\begin{aligned} F_q^p \left[\begin{matrix} (\alpha_1), (\alpha_2) \dots (\alpha_p) \\ (\beta_1), (\beta_2) \dots (\beta_q) \end{matrix} ; z \right] &= \frac{\Gamma(\beta_1) \Gamma(\beta_2) \dots \Gamma(\beta_q)}{\Gamma(\alpha_1) \Gamma(\alpha_2) \dots \Gamma(\alpha_p)} \\ & \Psi_q^p \left[\begin{matrix} (\alpha_1, 1), (\alpha_2, 1) \dots (\alpha_p, 1) \\ (\beta_1, 1), (\beta_2, 1) \dots (\beta_q, 1) \end{matrix} ; z \right] \end{aligned} \quad (2.2)$$

Lemma 1. *Let $\alpha, \beta, \eta \in C$. Then, there exists the relation*

(a) If $\operatorname{Re}(\alpha) > 0$ and $\operatorname{Re}(\sigma) > \max\{0, \operatorname{Re}(\beta - \eta)\}$, then

$$(I_{0^+}^{\alpha, \beta, \eta} t^{\sigma-1})(x) = \frac{\Gamma(\sigma)\Gamma(\sigma + \eta - \beta)}{\Gamma(\sigma - \beta)\Gamma(\sigma + \alpha - \eta)} x^{\sigma-\beta-1} \quad (2.3)$$

In particular, for $x > 0$, we have

$$(I_{0^+}^{\alpha} t^{\sigma-1})(x) = \frac{\Gamma(\sigma)}{\Gamma(\sigma + \alpha)} x^{\sigma+\beta+1}, \quad (2.4)$$

$$\operatorname{Re}(\alpha) > 0, \operatorname{Re}(\sigma) > 0 \quad (2.5)$$

$$(I_{\eta\alpha}^{\alpha} t^{\sigma-1})(x) = \frac{\Gamma(\sigma + \eta)}{\Gamma(\sigma + \alpha - \eta)} x^{\sigma-1}$$

$$\operatorname{Re}(\alpha) > 0, \operatorname{Re}(\sigma) > -\operatorname{Re}(\eta).$$

If $\operatorname{Re}(\alpha) > 0$ and $\operatorname{Re}(\sigma) < \min\{\operatorname{Re}(\beta), \operatorname{Re}(\eta)\}$, then

$$(I_{-}^{\alpha, \beta, \eta} t^{\sigma-1})(x) = \frac{\Gamma(1 - \sigma + \beta)\Gamma(1 - \sigma + \eta - \beta)}{\Gamma(\sigma - \sigma)\Gamma(1 - \sigma + \alpha + \beta + \eta)} x^{\sigma-\beta-1} \quad (2.6)$$

In particular, for $x > 0$, we have

$$(I_{-}^{\alpha} t^{\sigma-1})(x) = \frac{\Gamma(1 - \alpha + \sigma)}{\Gamma(1 - \sigma)} x^{\sigma-\alpha-1} \quad (2.7)$$

$$0 < \operatorname{Re}(\alpha) < 1 - \operatorname{Re}(\sigma).$$

$$(K_{\eta, \alpha}^{-} t^{\sigma-1})(x) = \frac{\Gamma(1 - \sigma + \eta)}{\Gamma(1 - \sigma + \alpha + \eta)} x^{\sigma-1} \quad (2.8)$$

$$0 < \operatorname{Re}(\alpha) < 1 + \operatorname{Re}(\sigma).$$

We begin the exposition of the main results by presenting the composition formulas of generalized fractional integrals, (1.1) and (1.2), involving the generalized Mathieu series in terms of the Hadamard product (1.14) of the generalized Mathieu series (1.13) and the Fox-Wright function (2.1). It is emphasized that the results presented here, which are for a seemingly complicated series, can reveal their involved properties via the series of the two known functions.

Theorem 1. Let $\alpha, \beta, \eta, \sigma \in C$ and $\rho > 0, \mu > 0, r \in R$ be such that

$\operatorname{Re}(\alpha) > 0$ and $\operatorname{Re}(\sigma) > \max\{0, \operatorname{Re}(\beta - \eta)\}$. Then, the following Saigo hypergeometric fractional integral $I_{0^+}^{\alpha, \beta, \eta}$ of $S_{\mu}\left(r; \left(\frac{t}{2}\right)^{\rho}\right)$ holds true:

$$\begin{aligned} \left(I_{0^+}^{\alpha, \beta, \eta} \left\{ \left(\frac{t}{2} \right)^{\sigma-1} S_{\mu} \left(r, \left(\frac{t}{2} \right)^{\rho} \right) \right\} \right) (x) &= x^{\sigma-\beta} S_{\mu} \left(r, \left(\frac{x}{2} \right)^{\rho} \right) \\ &* \Psi_2^3 \left[(1, 1), (\sigma + \rho, \rho), (\sigma + \eta - \beta + \rho, \rho); \left(\frac{x}{2} \right)^{\rho} \right] \end{aligned} \quad (2.9)$$

Proof. From the definition of $S_{\mu}\left(r; \left(\frac{t}{2}\right)^{\rho}\right)$

$$S_{\mu}\left(r; \left(\frac{t}{2}\right)^{\rho}\right) = \sum_{k=1}^{\infty} \frac{2n \left(\frac{t}{2}\right)^{\rho k}}{(k^2 + r^2)^{\mu+1}}.$$

Using the definitions (1.1) and (1.13), by changing the order of integration and applying the relation (2.3), we find that $x > 0$

$$\begin{aligned} I_{0^+}^{\alpha, \beta, \eta} \left\{ \sum_{k \geq 1} \frac{2k \left(\frac{t}{2}\right)^{\rho k + \sigma - 1}}{(k^2 + r^2)^{\mu+1}} \right\} &= \sum_{k=1}^{\infty} \frac{2k \left(\frac{1}{2}\right)^{\rho k + \sigma - 1}}{(k^2 + r^2)^{\mu+1}} (I_{0^+}^{\alpha, \beta, \eta} t^{\rho k + \sigma - 1})(x) \\ &= x^{\sigma-\beta-1} \sum_{k=1}^{\infty} \frac{2k \left(\frac{1}{2}\right)^{\rho k + \sigma - 1}}{(k^2 + r^2)^{\mu+1}} \frac{\Gamma(\sigma + \rho k) \Gamma(\sigma + \eta - \beta + \rho k)}{\Gamma(\sigma + \eta - \beta + \rho k) \Gamma(\sigma + \alpha + \eta - \beta + \rho k)} x^{\rho k} \end{aligned}$$

by applying the Hadamard product (1.14), which in the view of (1.14) and (2.1), yields the desired formula (2.9).

$$= x^{\sigma-\beta-1} S_{\mu} \left(r, \left(\frac{x}{2} \right)^{\rho} \right) * \Psi_2^3 \left[(1, 1), (\sigma + \rho, \rho), (\sigma + \eta - \beta + \rho, \rho); \left(\frac{x}{2} \right)^{\rho} \right]$$

This complete the prove of Theorem 1.

Theorem 2. Let $\alpha, \beta, \eta, \sigma \in C$ and $\rho > 0, \mu > 0, r \in R$ be such that $\operatorname{Re}(\alpha) > 0$ and $\operatorname{Re}(\sigma) > \max\{0, \operatorname{Re}(\beta - \eta)\}$. Then, the following Saigo

hypergeometric fractional integral $I_{0^+}^{\alpha, \beta, \eta}$ of $S_{\mu}\left(r, \left(\frac{1}{2t}\right)^{\rho}\right)$ holds true:

$$\left(I_{0^+}^{\alpha, \beta, \eta} \left\{ \left(\frac{t}{2}\right)^{\sigma-1} S_{\mu}\left(r, \left(\frac{t}{2t}\right)^{\rho}\right) \right\} \right) (x) = x^{\sigma-\rho-\beta-1} S_{\mu}\left(r, \left(\frac{1}{2x}\right)^{\rho}\right) * \Psi_2^3 \left[(1, 1), (1 - \sigma + \beta + \rho, \rho), (1 - \sigma + \eta + \rho, \rho); \left(\frac{1}{2x}\right)^{\rho} \right]. \tag{2,10}$$

Proof. From the definition of $S_{\mu}\left(r; \left(\frac{t}{2}\right)^{\rho}\right)$

$$S_{\mu}\left(r; \left(\frac{t}{2}\right)^{\rho}\right) = \sum_{k=1}^{\infty} \frac{2n\left(\frac{1}{2t}\right)^{\rho k}}{(k^2 + r^2)^{\mu+1}}$$

Using the definitions (1.1) and (1.13), by changing the order of integration and applying the relation (2.3), we find that

$$\begin{aligned} I_{0^+}^{\alpha, \beta, \eta} \left\{ \sum_{k=1}^{\infty} \frac{2k\left(\frac{t}{2t}\right)^{\rho k + \sigma - 1}}{(k^2 + r^2)^{\mu+1}} \right\} &= \sum_{k=1}^{\infty} \frac{2k\left(\frac{1}{2}\right)^{\rho k + \sigma - 1}}{(k^2 + r^2)^{\mu+1}} (I_{0^+}^{\alpha, \beta, \eta} t^{-\rho k + \sigma - 1})(x) \\ &= x^{\sigma-\rho-\beta-1} \sum_{k=1}^{\infty} \frac{2k\left(\frac{1}{2}\right)^{\rho k + \sigma - 1}}{(k^2 + r^2)^{\mu+1}} \frac{\Gamma(1 - \sigma + \beta + \rho, \rho), (1 - \sigma + \eta + \rho k)}{\Gamma(1 - \sigma + \rho k)\Gamma(1 - \sigma + \alpha + \beta + \eta + \rho k)} x^{-\rho k} \end{aligned}$$

by applying the Hadamard product (1.14), which in the view of (1.14) and (2.1), yields the desired formula (33).

$$= x^{\sigma-\rho-\beta-1} S_{\mu}\left(r, \left(\frac{x}{2}\right)^{\rho}\right) * \Psi_2^3 \left[(1, 1), (1 - \sigma + \beta + \rho, \rho), (1 - \sigma + \eta + \rho, \rho); \left(\frac{1}{2x}\right)^{\rho} \right]$$

This complete the prove of Theorem 2.7.

Further, we deduce the fractional integral formulas for the classical Riemann-Liouville and Erdélyi-Kober fractional integral and differential operators by letting $\beta = -\alpha$ and $\beta = 0$ respectively, which are asserted by Corollaries 1-4 below.

Corollary 3. Let $\alpha, \sigma \in C$ and $\rho > 0, \mu > 0, r \in R$ be such that $\operatorname{Re}(\alpha) > 0$ and $\operatorname{Re}(\sigma) > 0$. Then, the following Riemann-Liouville fractional integral $S_\mu \left(r; \left(\frac{t}{2} \right)^\rho \right)$ holds true:

$$\begin{aligned} & \left(I_{0^+}^\alpha \left\{ \left(\frac{t}{2} \right)^{\sigma-1} S_\mu \left(r, \left(\frac{t}{2} \right)^\rho \right) \right\} \right) (x) \\ &= x^{\sigma+\alpha+\beta-1} S_\mu \left(r, \left(\frac{x}{2} \right)^\rho \right) * \Psi_2^3 \left[\begin{matrix} (1,1), (\sigma+\rho, \rho) \\ (\sigma+\alpha+\rho\rho) \end{matrix}; \left(\frac{x}{2} \right)^\rho \right]. \end{aligned}$$

Corollary 4. Let $\alpha, \sigma \in C$ and $\rho > 0, \mu > 0, r \in R$ be such that $\operatorname{Re}(\alpha) > 0$ and $\operatorname{Re}(\sigma) > -\operatorname{Re}(\eta)$. Then, the following Erdélyi-Kober fractional integral $I_{\eta, \alpha}^+$ of $S_\mu \left(r; \left(\frac{t}{2} \right)^\rho \right)$ holds true:

$$\begin{aligned} & \left(I_{0^+}^\alpha \left\{ \left(\frac{t}{2} \right)^{\sigma-1} S_\mu \left(r, \left(\frac{t}{2} \right)^\rho \right) \right\} \right) (x) \\ &= x^{\sigma+\rho+\beta-1} S_\mu \left(r, \left(\frac{x}{2} \right)^\rho \right) * \Psi_2^3 \left[\begin{matrix} (1,1), (\sigma+\eta+\rho, \rho) \\ (\sigma+\alpha+\eta+\rho, \rho) \end{matrix}; \left(\frac{x}{2} \right)^\rho \right]. \end{aligned}$$

Corollary 5. Let $\alpha, \sigma \in C$ and $\rho > 0, \mu > 0, r \in R$ be such that $0 < \operatorname{Re}(\alpha) < 1 - \operatorname{Re}(\sigma)$. Then, the following Riemann-Liouville fractional integral I_-^α of $S_\mu \left(r; \left(\frac{1}{2t} \right)^\rho \right)$ holds true:

$$\begin{aligned} & \left(I_-^\alpha \left\{ \left(\frac{t}{2} \right)^{\sigma-1} S_\mu \left(r, \left(\frac{1}{2t} \right)^\rho \right) \right\} \right) (x) \\ &= x^{\sigma+\alpha-\rho-1} S_\mu \left(r, \left(\frac{1}{2x} \right)^\rho \right) * \Psi_1^2 \left[\begin{matrix} (1,1), (1-\sigma-\alpha+\rho, \rho) \\ (1-\sigma+\rho, \rho) \end{matrix}; \left(\frac{1}{2x} \right)^\rho \right]. \end{aligned}$$

Corollary 6. Let $\alpha, \sigma \in C$ and $\rho > 0, \mu > 0, r \in R$ be such that $0 < \operatorname{Re}(\alpha)$ and $\operatorname{Re}(\sigma) > 1 + \operatorname{Re}(\sigma)$. Then, the following Erdélyi-Kober

fractional integral I_-^α of $S_\mu\left(r; \left(\frac{1}{2t}\right)^\rho\right)$ holds true:

$$\begin{aligned} & K_{\eta, \alpha}^- \left(\left\{ \left(\frac{t}{2} \right)^{\sigma-1} S_\mu \left(r, \left(\frac{1}{2t} \right)^\rho \right) \right\} \right) (x) \\ &= x^{\sigma-\rho-1} S_\mu \left(r, \left(\frac{1}{2x} \right)^\rho \right) * \Psi_1^2 \left[\begin{matrix} (1, 1), (1 - \sigma - \alpha + \rho, \rho) \\ (1 - \sigma + \alpha + \eta + \rho, \rho) \end{matrix}; \left(\frac{1}{2x} \right)^\rho \right]. \end{aligned}$$

Acknowledgments

The authors thanks to the anonymous referees for valuable comments and useful suggestions, improving the quality of the article.

References

- [1] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, Theory and Applications of Fractional Differential Equations, North-Holland Mathematical Studies; Elsevier (North-Holland) Science Publishers: New York, NY, USA, Volume 204 2006.
- [2] V. S. Kiryakova, Generalized Fractional Calculus and Applications; Pitman Research Notes in Mathematics Series; Longman Scientific and Technical: Harlow, UK; John Wiley and Sons, Inc.: New York, NY, USA, 1993.
- [3] A. M. Mathai, R. K. Saxena and H. J. Haubold, The H-Functions: Theory and Applications; Springer: New York, NY, USA, 2010.
- [4] S. G. Samko, A. A. Kilbas and O. I. Marichev, Fractional Integrals and Derivatives: Theory and Applications; Translated from the Russian: Integrals and Derivatives of Fractional Order and Some of Their Applications (Nauka i Tekhnika, Minsk, 1987); Gordon and Breach Science Publishers: Reading, UK, 1993.
- [5] H. M. Srivastava, R. K. Saxena, Operators of fractional integration and their applications, Appl. Math. Comput. 118 (2001), 1-52.
- [6] M. Saigo, A remark on integral operators involving the Gauss hypergeometric functions, Math. Rep. Kyushu Univ. 11 (1978), 135-143.
- [7] E. L. Mathieu, Traité de Physique Mathématique, VI-VII: Théorie de L'élasticité des Corps Solides; Gauthier-Villars: Paris, France, 1890.
- [8] T. K. Pogány, H. M. Srivastava and Ž. Tomovski, Some families of Mathieu a-series and alternating Mathieu a-series, Appl. Math. Comput. 173 (2006), 69-108.
- [9] O. Emersleben, Über die Reihe $\sum_{k=1}^{\infty} \frac{k}{(k^2 + r^2)^2}$, Math. Ann. 125 (1952), 165-171.

- [10] P. Cerone, C. T. Lenard, On integral forms of generalized Mathieu series, *JIPAM J. Inequal, Pure Appl. Math.* 4 (2003), 1-11.
- [11] G. V. Milovanović and T. K. Pogány, New integral forms of generalized Mathieu series and related applications, *Appl. Anal. Discret. Math.* 7 (2013), 180-192.
- [12] P. H. Diananda, Some inequalities related to an inequality of Mathieu, *Math. Ann.* 250 (1980), 95-98.
- [13] H. M. Srivastava, Ž. Tomovski and D. Leškovski, Some families of Mathieu type series and Hurwitz-Lerch zeta functions and associated probability distributions, *Appl. Comput. Math.* 14 (2015), 349-380.
- [14] Ž. Tomovski and T. K. Pogány, Integral expressions for Mathieu-type power series and for the Butzer-Flocke-Hauss W-function, *Fract. Calc. Appl. Anal.* 14 (2011), 623-634.
- [15] Á. Baricz, P. L. Butzer and T. K. Pogány, Alternating Mathieu series, Hilbert-Eisenstein series and their generalized Omega functions, In *Analytic Number Theory, Approximation Theory, and Special Functions*; T. Rassias, G. V. Milovanović, Eds.; In Honor of Hari M. Srivastava; Springer: New York, NY, USA, (2014), 775-808.
- [16] J. Choi, R. K. Parmar and T. K. Pogány, Mathieu-type series built by (p, q) – Extended Gaussian hypergeometric function, *Bull. Korean Math. Soc.* 54 (2017), 789-797.
- [17] J. Choi and H. M. Srivastava, Mathieu series and associated sums involving the Zeta functions, *Comput. Math. Appl.* 59 (2010), 861-867.
- [18] N. Elezović, H. M. Srivastava and Ž. Tomovski, Integral representations and integral transforms of some families of Mathieu type series, *Integral Transforms Spec. Funct.* 19 (2008), 481-495.
- [19] T. K. Pogány and R. K. Parmar, On p-extended Mathieu series; submitted manuscript, arXiv2016, arXiv:1606.08369.
- [20] T. K. Pogány and Ž. Tomovski, Bounds improvement for alternating Mathieu type series, *J. Math. Inequal.* 4 (2010), 315-324.
- [21] R. K. Saxena and T. K. Pogány, R. Saxena, Integral transforms of the generalized Mathieu series, *J. Appl. Math. Stat. Inform.* 6 (2010), 5-16.
- [22] Ž. Tomovski, Integral representations of generalized Mathieu series via Mittag-Leffler type functions, *Fract. Calc. Appl. Anal.* 10 (2007), 127-138.
- [23] Ž. Tomovski, New integral and series representations of the generalized Mathieu series, *Appl. Anal. Discret. Math.* 2 (2008), 205-212.
- [24] Ž. Tomovski and T. K. Pogány, New upper bounds for Mathieu-type series, *Banach J. Math. Anal.* 3 (2009), 9-15.
- [25] V. Kiryakova, On two Saigo's fractional integral operators in the class of univalent functions, *Fract. Calc. Appl. Anal.* 9 (2006), 159-176.
- [26] T. Pohlen, The Hadamard Product and Universal Power Series, Ph.D. Thesis, Universität Trier, Trier, Germany, 2009.

- [27] H. M. Srivastava, R. Agarwal and S. Jain, Integral transform and fractional derivative formulas involving the extended generalized hypergeometric functions and probability distributions, *Math. Method Appl. Sci.* 40 (2017), 255-273.
- [28] R. Srivastava, R. Agarwal and S. Jain, A family of the incomplete hypergeometric functions and associated integral transform and fractional derivative formulas, *Filomat* 31 (2017), 125-140.
- [29] H. M. Srivastava and P. W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Ellis Horwood Ltd.: Chichester, UK, 1985.