



## CARTESIAN PRODUCTS OF FUZZY PRIME IDEAL

A. KIRUTHIKA, B. APSANA BEGAM, C. MAHENDRAN  
and N. VIJAYASEETHA

Department of Mathematics  
Dhanalakshmi Srinivasan College of Arts  
and Science For Women (Autonomous)  
Perambalur, India  
E-mail: akirushanmu1737@gmail.com

### Abstract

In this paper, the notation of fuzzy prime ideals and maximum prime ideals are introduced by applying the general theory of algebraic fuzzy system.

### Introduction

A fuzzy set is outlined mathematically by distribution to every doable individual within the universe of discourse a price representing its grade membership within the fuzzy set. This grade corresponds to the degree to that that individual is comparable or compatible with the construct diagrammatic by the fuzzy set. Thus individuals may belong to a greater or lesser degree as indicated by larger or smaller membership grade. These membership grades square measure fairly often diagrammatic by complex number values go within the interval between zero and one.

### Fuzzy Prime Ideals

**Definition.** Let  $\mu$  be any fuzzy ideal of a ring  $R$ ,  $t \in [0, 1]$  and  $t \leq \mu(0)$ . The ideal  $\mu_t$  is called a level ideal of  $\mu$ .

**Definition.** A fuzzy ideal  $\mu$  of a ring  $R$  is called prime, if the ideal  $\mu_t$ ,

---

2020 Mathematics Subject Classification: 03E72.

Keywords: Fuzzy Prime Ideals, Maximum Prime Ideals.

Received November 2, 2021; Accepted November 15, 2021

where  $\mu(0) = t$  is a prime ideal of  $R$ .

**Result**

If  $\mu$  is any fuzzy prime ideal of a ring  $R$ . Then  $\mu(xy) = \max(\mu(x), \mu(y))$ .

**Result**

If  $\mu$  and  $\theta$  are any fuzzy prime ideal of a ring  $R$ . Then  $\mu \cap \theta$  is a fuzzy prime ideal of  $R \leftrightarrow \mu \subseteq \theta$  (or)  $\theta \subseteq \mu$ .

**Theorem.**  $I \in I_m \mu$  and let  $\theta$  be any fuzzy prime ideal of  $R$ . Then  $\mu \cap \theta$  is a fuzzy prime ideal of the ring.

$$\mu_t = \{x \in R / \mu(x) = 1\}$$

**Proof.**  $\theta$  be any fuzzy prime ideal of  $R$ .

**Case (i).** If  $\theta$  is constant

Let  $\theta(r) = c$  for all  $r \in R$ . [since for all  $x \in \mu_t$ ]

Then, we have

$$\begin{aligned} (\mu \cap \theta)x &= \min(\mu(x), \theta(x)) \\ &= \min(1, c) \\ &= c \end{aligned}$$

Hence  $\mu \cap \theta$  is a fuzzy prime ideal of  $\mu_t$ .

**Case (ii).** If  $\theta$  is non-constant

Then there exists  $\alpha \in [0, 1]$  such that

$$\theta(x) = \begin{cases} 1 & \text{if } x \in \theta_t \\ \alpha & \text{if } x \in R \sim \theta_t \end{cases}$$

Where  $\theta_t = \{x \in R / \theta(x) = 1\}$

Also  $\theta_t$  is a prime ideal of  $R$

$\theta_t \cap \mu_t$  is a prime ideal of  $\mu_t$

$$(\mu \cap \theta) = \begin{cases} 1 & \text{if } x \in \theta_t \cap \mu_t \\ \alpha & \text{if } x \in \mu_t - (\theta_t \cap \mu_t) \end{cases}$$

Hence  $\mu \cap \theta$  is a fuzzy prime ideal of  $\mu_t$

Hence the proof.

**Theorem.** *If  $f$  is a homomorphism from a ring  $R$  onto a ring  $R'$  and  $\mu$  is any  $f$ -invariant fuzzy prime ideal of  $R$ . Then  $f(\mu)$  is fuzzy prime ideal of  $R'$ .*

(or)

$f(\mu)$  is a fuzzy prime ideal of  $R'$ . Prove that  $\sigma' \subseteq f(\mu)$  (or)  $\theta' \subseteq f(\mu)$ .

**Proof.** Let  $\sigma'$  and  $\theta'$  be any two fuzzy ideal of  $R'$  such that  $\sigma'\theta' \subseteq f(\mu)$

Then,

$$f^{-1}(\sigma'\theta') \subseteq f^{-1}(f(\mu)) = \mu \text{ [by result } f^{-1}(f(\mu)) = \mu \text{]}$$

$$f^{-1}(\sigma')f^{-1}(\theta') \subseteq \mu$$

Either,

$$\sigma' = f(f^{-1}(\sigma')) \subseteq f(\mu)$$

$$\sigma' \subseteq f(\mu) \text{ (or)}$$

$$\theta' = f(f^{-1}(\theta')) \subseteq f(\mu)$$

$$\theta' \subseteq f(\mu)$$

$$\sigma'\theta' \subseteq f(\mu)$$

Accordingly,

$f(\mu)$  is fuzzy prime ideal of  $R$ .

Hence the proof.

**Theorem.** *Let  $f$  be a homomorphism from a ring  $R$  onto a ring  $R'$  if  $\mu'$  and  $\theta'$  are any two fuzzy ideals of  $R'$  then the following holds,*

$$f^{-1}(\mu)f^{-1}(\theta') \subseteq f^{-1}(\mu'\theta')$$

**Proof.** Let  $x \in R$  and  $\varepsilon > 0$  be given the set

$$\alpha = (f^{-1}(\mu')f^{-1}(\theta'))(x) \text{ and}$$

$$\beta = (f^{-1}(\mu'\theta'))(x)$$

Then,

$$\alpha - \varepsilon < \sup(\min(f^{-1}(\mu'), (f^{-1}(\theta'))(x_1)))$$

$$x = x_1x_2$$

$$= \sup(\min(\mu'(f(x_1)), \theta'(f(x_1))))$$

$$x = x_1x_2$$

$$\alpha - \varepsilon < \min(\mu'(f(x_1)), \theta'(f(x_2))) \text{ for some } x = x_1x_2 \in R \text{ such that } x = x_1x_2$$

$$\leq (\mu'\theta')(f(x_1x_2))$$

$$= f^{-1}(\mu'\theta')(x) = \beta$$

$$\alpha - \varepsilon \leq \beta$$

Since  $\varepsilon$  is an arbitrary.

Hence,

$$f^{-1}(\mu')f^{-1}(\theta') \subseteq f^{-1}(\mu'\theta')$$

**Theorem.** *If  $f$  is an homomorphism from a ring  $R$  onto a ring  $R'$  and  $\mu'$  is any fuzzy prime ideal of  $R$ .*

**Proof.**  $\mu\sigma \leq f^{-1}(\mu)$

Then,

$$f(\mu\sigma) \subseteq f(f^{-1}(\mu)) = \mu'$$

By using theorem,

“Let  $f$  be a homomorphism from a  $r$  in  $R$  onto a ring and  $R'$  if  $\mu'$  and  $\theta'$  are any two fuzzy ideals of  $R'$  then the following holds,

$$f^{-1}(\mu')f^{-1}(\theta) \subseteq f^{-1}(\mu'\theta)$$

$$f(\mu)f(\sigma) \subseteq \mu'$$

Either  $f(\mu) \subseteq \mu'$  (or)

$$f(\sigma) \subseteq \mu'$$

Since  $\mu'$  is fuzzy prime,

Either,

$$f^{-1}(f(\mu)) \subseteq f^{-1}(\mu')$$

(or)

$$f^{-1}(f(\sigma)) \subseteq f^{-1}(\mu)$$

$$\mu \subseteq f^{-1}(\mu')$$

(or)

$$\sigma \subseteq f^{-1}(\mu')$$

Hence,

$f^{-1}(\mu)$  is fuzzy prime ideal of  $R$ .

**Theorem.** *If  $\mu$  and  $\sigma$  be fuzzy prime ideals of  $R$  then  $\mu \times \sigma$  is fuzzy prime ideal of  $R \times R$ .*

**Proof.** We know that the cartesian product of any two fuzzy ideals is fuzzy ideal by (Malik and Mordeson, 1991)

So it is enough to show that

$$\forall (a, b), (c, d) \in R \times R$$

Either,

$$\mu \times \sigma((a, b)(c, d)) = \mu \times \sigma((a, b))$$

$$\mu \times \sigma((a, b)(c, d)) = \mu \times \sigma((c, d))$$

Since  $\mu$  and  $\sigma$  is fuzzy prime ideals of  $R$  for all  $a, b, c, d \in R$

Either  $\mu(ac) = \mu(a)$  or else  $\mu(ac) = \mu(c)$

Either  $\mu(bd) = \mu(b)$  or else  $\mu(bd) = \mu(d)$

Then,

$$\begin{aligned} \mu \times \sigma((a, c)(b, d)) &= \mu \times \sigma(ac, bd) \\ &= \min(\mu(ac), \sigma(bd)) \\ &= \mu(ac) = \mu(a) = \mu \times \sigma(a, b) \\ &\text{or} \\ &= \mu(ac) = \mu(c) = \mu \times \sigma(c, d) \end{aligned}$$

Similarly,

$$\begin{aligned} \mu \times \sigma((a, c)(c, d)) &= \mu \times \sigma(ac, cd) \\ &= \min(\mu(ac), \sigma(cd)) \\ &= \sigma(cd) = \sigma(c) = \mu \times \sigma(a, c) \\ &\text{or} \\ &= \sigma(cd) = \sigma(d) = \mu \times \sigma(c, d) \end{aligned}$$

Therefore if  $\mu$  and  $\sigma$  are fuzzy prime ideals of  $R$  then  $\mu \times \sigma$  is a fuzzy prime ideal of  $R \times R$ .

**Theorem.** *If  $\mu$  and  $\sigma$  be fuzzy semi prime ideals of  $R$  then  $\mu \times \sigma$  is a fuzzy semi prime ideal of  $R \times R$ .*

**Proof.** We know that the cartesian product of any two fuzzy ideals is fuzzy ideal by (Malik and Mordeson, 1991).

So it is enough to show that for all  $(a, b) \in R \times R$ .

Either,

$$\mu \times \sigma((a, b)^n) = \mu \times \sigma(a, b)$$

Since  $\mu$  and  $\sigma$  is fuzzy semi prime ideals of  $R$  for all  $a, b \in R, \forall n \in \mathbb{N}^+$ .

$$\mu(a)^n = \mu(a) \text{ and } \sigma(b)^n = \mu(b)$$

Then,

$$\begin{aligned} \mu \times \sigma((a, b)^n) &= \mu \times \sigma(a^n, b^n) \\ &= \min (\mu(a^n), \sigma(b^n)) \\ &= \mu(a^n) \\ &= \mu(a) \\ &= \mu \times \sigma(a, b) \end{aligned}$$

$$\begin{aligned} \mu \times \sigma((a, b)^n) &= \mu \times \sigma(a^n, b^n) \\ &= \min (\mu(a^n), \sigma(b^n)) \\ &= \sigma(b^n) \\ &= \sigma(b) \\ &= \mu \times \sigma(a, b) \end{aligned}$$

Therefore  $\mu \times \sigma$  is a fuzzy semi prime ideal of  $R \times R$ .

**Theorem.** *If  $\mu$  and  $\sigma$  be fuzzy semi primary ideals of  $R$  then  $\mu \times \sigma$  is a fuzzy semi primary ideal of  $R \times R$ .*

**Proof.** Since,  $\mu$  and  $\sigma$  be fuzzy semi primary ideals of  $R$  then for all  $a, b, c, d \in R$ .

Either  $\mu(ab) \leq \mu(a^n)$  or else  $\mu(ab) \leq \mu(b^m)$  for some  $m, n \in \mathbb{N}^+$ .

Either  $\sigma(cd) \leq \sigma(d^k)$  or else  $\sigma(cd) \leq \sigma(c^l)$  for some  $k, l \in \mathbb{N}^+$ .

Let  $t = \max(m, n)$  and

$$s = \max(k, l)$$

Then,

$$\begin{aligned}
 \mu \times \sigma((a, c)(b, d)) &= \mu \times \sigma(ab, cd) \\
 &= \min(\mu(ab), \sigma(cd)) \\
 &= \mu(ab) \\
 &\leq \mu(a^t) \\
 &= \mu \times \sigma((a, c)^t)
 \end{aligned}$$

Or else,

$$\begin{aligned}
 \mu \times \sigma((a, c)(b, d)) &= \mu \times \sigma(ab, cd) \\
 &= \min(\mu(ab), \sigma(cd)) \\
 &= \sigma(cd) \\
 &\leq \sigma(d^s) \\
 &= \mu \times \sigma((b, d)^s)
 \end{aligned}$$

Therefore  $\mu \times \sigma$  is fuzzy semi primary ideal of  $R \times R$ .

Hence the proof.

**Corollary.** *A fuzzy ideal  $\mu \times \sigma$  of  $R \times R$  is fuzzy semi primary ideal if and only if the level ideals  $t \in I_m \mu \times \sigma, (\mu \times \sigma)_t$  are semi primary ideals of  $R \times R$ s.*

### Conclusion

The study of semi prime ideals in universal algebras is under investigation by the authors using the commentator of fuzzy ideals. Moreover, the radical of fuzzy ideals in universal algebra will be studied and would be applied to characterize the central properties of fuzzy semi prime ideals.



**References**

- [1] A. R. Meenakshi, Fuzzy algebra, MJP Publisher (2019), 38-51.
- [2] A. R. Meenakshi, Fuzzy matrix Theory and Applications, MJP Publisher (2019), 89-102.
- [3] Rajesh Kumar, University of Delhi Press, Fuzzy algebra, Publication Division (1993), 66-78.
- [4] Dr. W. B. Vasantha Kandasamy, Fuzzy Algebra, Indian Institute of Technology of Chennai, American Research Press (2003), 39-63.
- [5] Dr. W. B. Vasantha Kandasamy, Smarandache Fuzzy Algebra, American Research Press, Rehoboth (2003), 333-340.